

On the I-loop effective action of string compactifications

Michael Haack, (LMU Munich)
Zlatibor, September 8, 2018

I407.0027 (with Marcus Berg, Jin U Kang and Stefan Sjörs)
I511.03957 (with Jin U Kang)
+ I805.00817 (with Jin U Kang)

Overview

- Motivation
- Calculational setup
- Results

Motivation

(Perturbative) quantum corrections to effective action can be important

- if “zero effect” at tree level
(e.g. no-scale structure of potential)
- increase possibilities for model building and moduli stabilization, cf. “Large Volume Scenario”
[Balasubramanian, Berglund, Conlon, Quevedo]

$\mathcal{N} = 1, d = 4$ Supergravity

$$\begin{aligned}\frac{\mathcal{L}_{\text{bos}}}{(-g)^{1/2}} = & \frac{1}{2\kappa^2} R - K_{,\bar{I}J} D_\mu \bar{\Phi}^I D^\mu \Phi^J - \frac{1}{4} \text{Re}(f_{ab}(\Phi)) F_{\mu\nu}^a F^{b\mu\nu} \\ & - \frac{1}{8} \text{Im}(f_{ab}(\Phi)) \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu}^a F_{\rho\sigma}^b - V(\Phi, \bar{\Phi})\end{aligned}$$

with $V(\Phi, \bar{\Phi}) = e^K (G^{\bar{I}J} D_{\bar{I}} \bar{W} D_J W - 3|W|^2) + \text{Re}(f_{ab}) \mathcal{D}^a \mathcal{D}^b$


$$D_J W = \partial_{\Phi^J} W + (\partial_{\Phi^J} K) W$$

- Kähler potential K
- Gauge kinetic function f_{ab}
- Superpotential W

Quantum Corrections

- Superpotential $W = W^{\text{tree}} + W^{\text{non-pert}}$
- Gauge kinetic function $f = f^{\text{tree}} + f^{\text{1-loop}} + f^{\text{non-pert}}$
- Kähler potential $K = K^{\text{tree}} + \sum_{n=1}^{\infty} K^{n\text{-loop}} + K^{\text{non-pert}}$

String perturbation theory



Goal and Method

- Goal: Calculate string 1-loop corrections to Kähler potential K of moduli fields in type I theory

In applications this would give you direct access to corrections to the potential V

- Method: 1-loop scattering amplitudes in type I

Complications:

- Result from string amplitudes not in Einstein frame, but

$$S = \frac{1}{\kappa^2} \int d^4x \sqrt{-g} [(1 + \delta E) \frac{1}{2} R + \text{kinetic terms}]$$

Need Weylrescaling: $g_{\mu\nu}^{(E)} = \underbrace{(1 + \delta E) g_{\mu\nu}}_{\equiv \Omega^{-2}}$

⇒ kinetic terms multiplied by Ω^2

- String theory naturally calculates

$$\tilde{G}_{\bar{I}J} \partial_\mu \bar{\varphi}^I \partial^\mu \varphi^J = \left(G_{\bar{I}J}^{(0)}(\varphi) + \tilde{G}_{\bar{I}J}^{(1)}(\varphi) \right) \partial_\mu \bar{\varphi}^I \partial^\mu \varphi^J$$

with $\Phi^I = \Phi^I(\varphi)$

Complications:

- Result from string amplitudes not in Einstein frame, but

$$S = \frac{1}{\kappa^2} \int d^4x \sqrt{-g} [(1 + \delta E) \frac{1}{2} R + \text{kinetic terms}]$$

Need Weylrescaling: $g_{\mu\nu}^{(E)} = \underbrace{(1 + \delta E) g_{\mu\nu}}_{\equiv \Omega^{-2}}$

\Rightarrow kinetic terms multiplied by Ω^2

- String theory naturally calculates

$$\tilde{G}_{\bar{I}J} \partial_\mu \bar{\varphi}^I \partial^\mu \varphi^J = \left(G_{\bar{I}J}^{(0)}(\varphi) + \tilde{G}_{\bar{I}J}^{(1)}(\varphi) \right) \partial_\mu \bar{\varphi}^I \partial^\mu \varphi^J$$

with $\Phi^I = \Phi^I(\varphi)$ (e.g. $T = c + i\tau$)
RR field $e^{-\Phi_{10}} \mathcal{V}$

- Thus, string theory gives you (after Weyl-rescaling):

$$G_{\bar{I}J} \partial_\mu \bar{\varphi}^I \partial^\mu \varphi^J = \left(G_{\bar{I}J}^{(0)}(\varphi) + \underbrace{G_{\bar{I}J}^{(1)}(\varphi)}_{\tilde{G}_{\bar{I}J}^{(1)}(\varphi) - G_{\bar{I}J}^{(0)}(\varphi)\delta E(\varphi)} \right) \partial_\mu \bar{\varphi}^I \partial^\mu \varphi^J$$

- Suppose $G_{\bar{I}J}^{(0)}(\varphi) \partial_\mu \bar{\varphi}^I \partial^\mu \varphi^J = \frac{\partial^2 K^{(0)}(\Phi^{(0)})}{\partial \bar{\Phi}^{(0)I} \partial \Phi^{(0)J}} \partial_\mu \bar{\Phi}^{(0)I} \partial^\mu \Phi^{(0)J}$ (★)

then in general $G_{\bar{I}J}^{(1)}(\varphi) \partial_\mu \bar{\varphi}^I \partial^\mu \varphi^J \neq \frac{\partial^2 K^{(1)}(\Phi^{(0)})}{\partial \bar{\Phi}^{(0)I} \partial \Phi^{(0)J}} \partial_\mu \bar{\Phi}^{(0)I} \partial^\mu \Phi^{(0)J}$

- Solution: $\Phi^I = \Phi^I(\varphi) = \Phi^{(0)I}(\varphi) + \Phi^{(1)I}(\varphi)$

⇒ Get additional contributions to the 1-loop metric from inserting $\Phi^{(0)I} = \Phi^I - \Phi^{(1)I}$ into (★) such that

$$\left(G_{\bar{I}J}^{(0)}(\varphi) + G_{\bar{I}J}^{(1)}(\varphi) \right) \partial_\mu \bar{\varphi}^I \partial^\mu \varphi^J = \frac{\partial^2 \left(K^{(0)}(\Phi) + K^{(1)}(\Phi) \right)}{\partial \bar{\Phi}^I \partial \Phi^J} \partial_\mu \bar{\Phi}^I \partial^\mu \Phi^J$$

\implies read off $K^{(1)}$

- Upshot: Need 1-loop corrections to

- (i) scalar metric $\tilde{G}_{\bar{I}J}^{(1)}(\varphi)$
- (ii) Einstein-Hilbert term δE
- (iii) definition of field variables $\Phi^{(1)I}(\varphi)$




Calculational setup

Some generalities of the amplitude calculations:


- Aim: read off scalar metric from scalar 2-pt fct.
- 2-pt fct. = 0 on-shell
- Trick: use $p_1 + p_2 \neq 0 \iff \lambda \equiv p_1 \cdot p_2 \neq 0$
in intermediate steps [Atick, Dixon, Sen; Minahan; Antoniadis, Bachas, Fabre, Partouche, Taylor; Antoniadis, Kirtsis, Rizos; cf. also Kirlitsis, Kounnas, ...]
- $\langle \varphi_i \varphi_j \rangle = \lambda G_{ij} + \mathcal{O}(\lambda^2)$
- Similarly for gravitons: $\langle hh \rangle \sim \delta E p_2^\mu \epsilon_{1\mu\nu} \eta^{\nu\sigma} \epsilon_{2\sigma\rho} p_1^\rho$

$$T^6/\mathbb{Z}'_6$$

- $\Theta Z^1 = e^{2\pi i v_1} Z^1 \quad \Theta Z^2 = e^{2\pi i v_2} Z^2 \quad \Theta Z^3 = e^{2\pi i v_3} Z^3$
 $(v_1, v_2, v_3) = \left(\frac{1}{6}, -\frac{1}{2}, \frac{1}{3}\right)$



- Example \mathbb{Z}_3 :




- *Orientifold*: Identify under worldsheet reflection

$$\Omega | \circlearrowright \rangle = | \circlearrowleft \rangle$$


$$\frac{1 + \Omega}{2} | \circlearrowright \rangle \quad \text{unoriented eigenstate}$$

- Resulting 4D effective action has $\mathcal{N} = 1$
- Model contains D9- and D5-branes (wrapped around 3rd torus)
- In addition to torus, at 1-loop need to calculate:


Annulus:



Möbius strip:



Klein bottle:



All of them have Euler number $\chi = 0$

- E.g. annulus:

$$\mathcal{A} \sim \int_0^\infty \frac{dt}{t} \text{Tr}_{open} \left(\left[\frac{1}{6} \sum_{k=0}^5 \Theta^k \right] q^{(p^2+m^2)/2} VV \right)$$

$$= \frac{1}{6} \sum_{k=0}^5 \int_0^\infty \frac{dt}{t} \text{Tr}_{open} \left(\Theta^k q^{(p^2+m^2)/2} VV \right)$$

$e^{-\pi t}$

projection operator
on orbifold invariant states


vertex operator

- $\mathcal{N} = 1$ contribution, if strings twisted along all 3 tori

Results: Scalar kinetic term

Concretely consider:

$$\tau = \text{Im}(T_3) \text{ with } \tau^{(0)} \sim e^{-\Phi} \mathcal{V}_3$$

10D dilaton 
volume of 3rd torus measured
with string frame metric 

Note: $e^{2\Phi} = e^{2\Phi_4} \mathcal{V}$

4D dilaton 
overall volume measured
with string frame metric 

Results: Scalar kinetic term

Prior results:

- No contribution from disk-level [Lüst, Mayr, Richter, Stieberger]
- Sphere, torus and $\mathcal{N} = 2$ -sectors:

$$G^{(0)} = -\frac{1}{4(\tau^{(0)})^2} \left(1 + \frac{\zeta(3)\chi}{\mathcal{V}} \right)$$

$$\tilde{G}^{(1)} \sim e^{2\Phi_4} \left(\frac{\chi}{(\tau^{(0)})^2} + a_1 \frac{1}{(\tau^{(0)})^2 \mathcal{V}_3} E_2(U_3) + a_2 \frac{\mathcal{V}_2}{(\tau^{(0)})^2} E_2(-1/U_2) \right)$$

complex structure of 3rd torus

$$E_2(U) \equiv \sum_{(m,n) \neq (0,0)} \frac{(\text{Im}(U))^2}{|m + nU|^4}$$

Euler number (48 for \mathbb{Z}'_6)

[Antoniadis, Ferrara, Minasian, Narain;
Becker, Becker, M.H., Louis;
Berg, M.H., Körs]

- New result: $\mathcal{N} = 1$ sectors [Berg, M.H., Kang, Sjörs]
- Usual lore: $\mathcal{N} = 1$ sectors less interesting, because they do not lead to moduli dependent results
- Moduli dependence in $\mathcal{N} = 1$ sectors via:
 - ★ normalization of vertex operators
 - ★ dilaton factor in Einstein frame
- Expect further moduli dependence in $\mathcal{N} = 1$ in presence of world volume fluxes or for branes at angles

- For $\tau = \text{Im}(T_3)$ in \mathbb{Z}'_6 from $\mathcal{A}, \mathcal{M}, \mathcal{K}$: [Berg, M.H., Kang, Sjörs]

$$\tilde{G}_{(\mathcal{N}=1)}^{(1)} = e^{2\Phi_4} \frac{5}{2^9 \pi^3} \overbrace{\text{Cl}_2\left(\frac{\pi}{3}\right)}^{\approx 3.197 \cdot 10^{-4}} \frac{1}{(\tau^{(0)})^2}$$

2nd Clausen function $\text{Cl}_2(\varphi) = \sum_{k=1}^{\infty} \frac{\sin(k\varphi)}{k^2}$

- This is a correction to the usual torus contribution

$$\sim e^{2\Phi_4} \frac{\chi}{(\tau^{(0)})^2}$$

Results: EH-term

- Prior results:

★ Type II: [Antoniadis, Ferrara, Minasian, Narain]

$$S^{(II)} = \frac{1}{\kappa^2} \int d^4x \sqrt{-g} \left[e^{-2\Phi_4} + \chi \left(\zeta(3) \frac{e^{-2\Phi_4}}{\mathcal{V}} \pm \frac{\pi^2}{6} \right) \right] \frac{R}{2}$$

IIB

IIA

★ Type I (on $K_3 \times T^2$): [Antoniadis, Bachas, Fabre, Partouche, Taylor]

$$S^{(I)} = \frac{1}{\kappa^2} \int d^4x \sqrt{-g} \left[e^{-2\Phi_4} + a \underbrace{\frac{1}{\mathcal{V}_{T^2}} E_2(U)}_{\text{from } \mathcal{A}, \mathcal{M}, \mathcal{K}} \right] \frac{R}{2} + \dots$$

★ Heterotic string: No 1-loop contribution, i.e.

$$S^{(het)} = \frac{1}{\kappa^2} \int d^4x \sqrt{-g} \left[e^{-2\Phi_4} \left(1 + \frac{\chi\zeta(3)}{\mathcal{V}} \right) \right] \frac{R}{2} + \dots$$

[Antoniadis, Gava, Narain; Kirlitsis, Kounnas]

This can be understood via

- World sheet calculation: integrand of torus & higher loop graviton 2-point function is total derivative

[Kirlitsis, Kounnas, Petropoulos, Rizos]

- $10D$ R^4 -terms

- Type II: [Gross, Witten; Green, Schwarz; Grisaru, van de Ven, Zanon]

$$(\zeta(3)e^{-2\Phi} + \frac{\pi^2}{6})t_8 t_8 R^4 - (\zeta(3)e^{-2\Phi} \pm \frac{\pi^2}{6})\epsilon_{10} \epsilon_{10} R^4$$

leads to correction to 4D kinetic terms of scalars

leads to correction to 4D EH-term

- Heterotic: [Cai, Nunez; Gross, Sloan; Sakai, Tanii; Abe, Kubota, Sakai]

$$(\zeta(3)e^{-2\Phi} + \frac{\pi^2}{6})t_8 t_8 R^4 - \zeta(3)e^{-2\Phi} \epsilon_{10} \epsilon_{10} R^4$$

- How is this compatible with heterotic / type I duality?

[Tseytlin; Green, Rudra]

$$g_{\mu\nu}^{(I)} = e^{-\Phi_{het}} g_{\mu\nu}^{(het)} \quad , \quad \Phi_I = -\Phi_{het}$$

- How is this compatible with heterotic / type I duality?

[Tseytlin; Green, Rudra]

$$g_{\mu\nu}^{(I)} = e^{-\Phi_{het}} g_{\mu\nu}^{(het)} \quad , \quad \Phi_I = -\Phi_{het}$$

- Possible answer: $S^{(het)}$ in 10D contains

[Green, Rudra]

$$\sqrt{g^{(het)}} e^{-\Phi_{het}/2} J_0 E_{3/2}(e^{-\Phi_{het}}) - \sqrt{g^{(het)}} \frac{\pi^2}{6} \mathcal{I}_2$$

★ $J_0 = t_8 t_8 R^4 - \frac{1}{8} \epsilon_{10} \epsilon_{10} R^4$

★ $\mathcal{I}_2 = -\frac{1}{8} \epsilon_{10} \epsilon_{10} R^4 + \dots$

★ $E_{3/2} = \zeta(3) e^{-3/2 \Phi_{het}} + \frac{\pi^2}{6} e^{\Phi_{het}/2} + \text{non-pert.}$

S-duality invariant

- How is this compatible with heterotic / type I duality?

[Tseytlin; Green, Rudra]

$$g_{\mu\nu}^{(I)} = e^{-\Phi_{het}} g_{\mu\nu}^{(het)} \quad , \quad \Phi_I = -\Phi_{het}$$

- Possible answer: $S^{(het)}$ in 10D contains

[Green, Rudra]

$$\underbrace{\sqrt{g^{(het)}} e^{-\Phi_{het}/2} J_0 E_{3/2}(e^{-\Phi_{het}}) - \sqrt{g^{(het)}} \frac{\pi^2}{6} \mathcal{I}_2}_{\rightarrow \sqrt{g^{(I)}} e^{-\Phi_I/2} J_0 E_{3/2}(e^{-\Phi_I})}$$

★ $J_0 = t_8 t_8 R^4 - \frac{1}{8} \epsilon_{10} \epsilon_{10} R^4$

★ $\mathcal{I}_2 = -\frac{1}{8} \epsilon_{10} \epsilon_{10} R^4 + \dots$

★ $E_{3/2} = \zeta(3) e^{-3/2 \Phi_{het}} + \frac{\pi^2}{6} e^{\Phi_{het}/2} + \text{non-pert.}$

S-duality invariant

- How is this compatible with heterotic / type I duality?

[Tseytlin; Green, Rudra]

$$g_{\mu\nu}^{(I)} = e^{-\Phi_{het}} g_{\mu\nu}^{(het)} \quad , \quad \Phi_I = -\Phi_{het}$$

- Possible answer: $S^{(het)}$ in 10D contains

[Green, Rudra]

$$\underbrace{\sqrt{g^{(het)}} e^{-\Phi_{het}/2} J_0 E_{3/2}(e^{-\Phi_{het}})}_{\rightarrow \sqrt{g^{(I)}} e^{-\Phi_I/2} J_0 E_{3/2}(e^{-\Phi_I})} - \underbrace{\sqrt{g^{(het)}} \frac{\pi^2}{6} \mathcal{I}_2}_{\rightarrow \sqrt{g^{(I)}} \frac{\pi^2}{6} e^{-\Phi_I} \mathcal{I}_2}$$

★ $J_0 = t_8 t_8 R^4 - \frac{1}{8} \epsilon_{10} \epsilon_{10} R^4$

★ $\mathcal{I}_2 = -\frac{1}{8} \epsilon_{10} \epsilon_{10} R^4 + \dots$

★ $E_{3/2} = \zeta(3) e^{-3/2 \Phi_{het}} + \frac{\pi^2}{6} e^{\Phi_{het}/2} + \text{non-pert.}$

S-duality invariant

- How is this compatible with heterotic / type I duality?

[Tseytlin; Green, Rudra]

$$g_{\mu\nu}^{(I)} = e^{-\Phi_{het}} g_{\mu\nu}^{(het)} \quad , \quad \Phi_I = -\Phi_{het}$$

- Possible answer: $S^{(het)}$ in 10D contains

[Green, Rudra]

$$\underbrace{\sqrt{g^{(het)}} e^{-\Phi_{het}/2} J_0 E_{3/2}(e^{-\Phi_{het}})}_{\rightarrow \sqrt{g^{(I)}} e^{-\Phi_I/2} J_0 E_{3/2}(e^{-\Phi_I})} - \underbrace{\sqrt{g^{(het)}} \frac{\pi^2}{6} \mathcal{I}_2}_{\rightarrow \sqrt{g^{(I)}} \frac{\pi^2}{6} e^{-\Phi_I} \mathcal{I}_2}$$

★ $J_0 = t_8 t_8 R^4 - \frac{1}{8} \epsilon_{10} \epsilon_{10} R^4$

★ $\mathcal{I}_2 = -\frac{1}{8} \epsilon_{10} \epsilon_{10} R^4 + \dots$

★ $E_{3/2} = \zeta(3) e^{-3/2 \Phi_{het}} + \frac{\pi^2}{6} e^{\Phi_{het}/2} + \text{non-pert.}$

S-duality invariant

\circledcirc Disk level!

- How is this compatible with heterotic / type I duality?

[Tseytlin; Green, Rudra]

$$g_{\mu\nu}^{(I)} = e^{-\Phi_{het}} g_{\mu\nu}^{(het)} \quad , \quad \Phi_I = -\Phi_{het}$$

- Possible answer: $S^{(het)}$ in 10D contains

[Green, Rudra]

$$\underbrace{\sqrt{g^{(het)}} e^{-\Phi_{het}/2} J_0 E_{3/2}(e^{-\Phi_{het}}) - \sqrt{g^{(het)}} \frac{\pi^2}{6} \mathcal{I}_2}_{\rightarrow \sqrt{g^{(I)}} e^{-\Phi_I/2} J_0 E_{3/2}(e^{-\Phi_I})} \quad \rightarrow \sqrt{g^{(I)}} \frac{\pi^2}{6} e^{-\Phi_I} \mathcal{I}_2$$

★ $J_0 = t_8 t_8 R^4 - \frac{1}{8} \epsilon_{10} \epsilon_{10} R^4$

★ $\mathcal{I}_2 = -\frac{1}{8} \epsilon_{10} \epsilon_{10} R^4 + \dots$

★ $E_{3/2} = \zeta(3) e^{-3/2 \Phi_{het}} + \frac{\pi^2}{6} e^{\Phi_{het}/2} + \text{non-pert.}$

Disk level!

Disk level correction
to 4D EH-term?

S-duality invariant

- New results for \mathbb{Z}'_6 : [M.H., Kang]
from $\mathcal{N} = 1$ -sectors of $\mathcal{A}, \mathcal{M}, \mathcal{K}$

$$\delta E = \frac{\chi}{(2\pi)^3} \left(2\zeta(3) + \frac{\pi^2}{3} e^{2\Phi_4} \right) + e^{2\Phi_4} \frac{5}{64\pi^2} \text{Cl}_2 \left(\frac{\pi}{3} \right)$$

$$- e^{2\Phi_4} \frac{5}{256\pi^2} \left[\frac{64\pi^2\alpha'}{\mathcal{V}_3} E_2(U_3) - \frac{12\pi^2\alpha'}{\mathcal{V}_2} E_2(U_2) - \frac{3\mathcal{V}_2}{4\pi^2\alpha'} E_2(-1/U_2) \right]$$

from $\mathcal{N} = 2$ -sectors of $\mathcal{A}, \mathcal{M}, \mathcal{K}$

- Follows closely a calculation by [Epple]
- Generalization to \mathbb{Z}_3 [M.H., Kang] and $\mathbb{Z}_6, \mathbb{Z}_7, \mathbb{Z}_{12}$
[with Tailin Li, unpublished]

Results: Field redefinitions

- Examples:

★ Type I (on $K_3 \times T^2$): $\tau = \tau^{(0)} + A^2 + \frac{E_2(U)}{e^{-\Phi}\mathcal{V}}$

[Antoniadis, Bachas, Fabre,
Partouche, Taylor]

open string scalars (cf. inflation in
string theory)

★ Type II on CY:

[Antoniadis, Minasian,
Theisen, Vanhove]

$$e^{-2\tilde{\Phi}_4} = e^{-2\Phi_4} \left(1 + a \frac{\chi}{\mathcal{V}} + \dots \right)$$

$$\tilde{\mathcal{V}} = \mathcal{V} \left(1 - 12 \frac{\zeta(2)}{(2\pi)^3} \chi e^{2\Phi_4} + \dots \right)$$

Determine form of *1-loop* field redefinitions using:

- ★ Kählerness of metric
- ★ Shift symmetries
- ★ Ansatz for 1-loop correction of metric
(dependence on volume moduli and dilaton)

- Tree level coordinates:

$$\star \quad T_j^{(0)} = c_j^{(0)} + i\tau_j^{(0)}, \quad j = 1, 2, 3$$

↗ component of
 RR 2-form C_2
 along j -th torus

$e^{-\Phi}\mathcal{V}_j$ ↗ volume of j -th torus
 (in string frame metric)

$$\star \quad T_0^{(0)} = c_0^{(0)} + i\tau_0^{(0)}$$

↗ dual to $C_{\mu\nu}$

$e^{-\Phi}\mathcal{V}$ ↗ overall volume

- Tree level Kähler potential:

$$K = - \sum_{I=0}^3 \ln \left(T_I^{(0)} - \bar{T}_I^{(0)} \right) \Rightarrow G_{I\bar{J}} \sim \frac{\delta_{IJ}}{(\tau_I^{(0)})^2}$$

- In principle one can calculate 1-loop corrections to metric using well-known vertex operators for \mathcal{V}_j , Φ_4 and RR-fields
- Express these (after Weyl-rescaling to Einstein frame) in terms of $c_I^{(0)}$, $\tau_I^{(0)}$ via

$$e^{2\Phi_4} = \frac{1}{\sqrt{\prod_{I=0}^3 \tau_I^{(0)}}} \quad , \quad \mathcal{V}_j = \tau_j^{(0)} \sqrt{\frac{\tau_0^{(0)}}{\prod_{i=1}^3 \tau_i^{(0)}}}$$

- Result:

$$\mathcal{L} = G_{c_I^{(0)} c_J^{(0)}}(\tau^{(0)}) \partial_\mu c_I^{(0)} \partial^\mu c_J^{(0)} + G_{\tau_I^{(0)} \tau_J^{(0)}}(\tau^{(0)}) \partial_\mu \tau_I^{(0)} \partial^\mu \tau_J^{(0)}$$

G independent of $c^{(0)}$ due to perturbative shift symmetry

$$\mathcal{L} = G_{c_I^{(0)} c_J^{(0)}}(\tau^{(0)}) \partial_\mu c_I^{(0)} \partial^\mu c_J^{(0)} + G_{\tau_I^{(0)} \tau_J^{(0)}}(\tau^{(0)}) \partial_\mu \tau_I^{(0)} \partial^\mu \tau_J^{(0)} \quad (\star)$$

- Aim: Find variables

$$T_I = c_I^{(0)} + c_I^{(1)}(c^{(0)}, \tau^{(0)}) + i\left(\tau_I^{(0)} + \tau_I^{(1)}(c^{(0)}, \tau^{(0)})\right)$$

such that (\star) becomes

$$\mathcal{L} = K_{I\bar{J}} \partial_\mu T_I \partial^\mu \bar{T}_{\bar{J}}$$

- Shift symmetry of c , i.e. $K = K(\tau)$, implies:

$$T = c + i\tau$$



★ $K_{T\bar{T}} \partial_\mu T \partial^\mu \bar{T} = \frac{1}{4} K_{\tau\tau} (\partial_\mu c \partial^\mu c + \partial_\mu \tau \partial^\mu \tau)$

(i) $G_{c_i \tau_j} = 0$

(ii) $G_{\tau_i \tau_j} = \frac{1}{4} \frac{\partial^2 K}{\partial \tau_i \partial \tau_j} = G_{c_i c_j}$

(iii) $\partial_{\tau_k} G_{c_i c_j} = \partial_{\tau_i} G_{c_j c_k} = \partial_{\tau_j} G_{c_k c_i}$

★ $c_I^{(1)} = 0$

Ansatz for 1-loop correction of metric: (motivated by string amplitudes)

- $\mathcal{N} = 1 :$
$$G_{\tau_I^{(0)} \tau_J^{(0)}}^{(1)} = \alpha_{IJ} \frac{e^{2\Phi_4}}{\tau_I^{(0)} \tau_J^{(0)}}$$
- $\mathcal{N} = 2 :$
$$G_{\tau_I^{(0)} \tau_J^{(0)}}^{(1)} = \frac{e^{2\Phi_4}}{\tau_I^{(0)} \tau_J^{(0)}} \sum_{\mathcal{N}=2-\text{sectors}} \alpha_{IJ}^{(l,n)}(U_l) \mathcal{V}_l^n$$

(similarly for $G_{c_I^{(0)} c_J^{(0)}}^{(1)}$)

$n = -1 :$ Closed string winding state exchange

$n = +1 :$ Closed string KK state exchange

- E.g. (for Z'_6) [M.H., Kang]

$\mathcal{T}, \mathcal{A}, \mathcal{M}, \mathcal{K}$

$\mathcal{N} = 1$

$$\delta\tau_3 = a_1 \sqrt{\frac{T_3^{(0)} - \bar{T}_3^{(0)}}{(T_0^{(0)} - \bar{T}_0^{(0)})(T_1^{(0)} - \bar{T}_1^{(0)})(T_2^{(0)} - \bar{T}_2^{(0)})}}$$

$$+ a_2 \frac{(T_3^{(0)} - \bar{T}_3^{(0)})E_2(U_2)}{(T_0^{(0)} - \bar{T}_0^{(0)})(T_2^{(0)} - \bar{T}_2^{(0)})}$$

$$+ a_3 \frac{E_2(-1/U_2)}{(T_1^{(0)} - \bar{T}_1^{(0)})} + a_4 \frac{E_2(U_3)}{(T_0^{(0)} - \bar{T}_0^{(0)})}$$

analog of correction by
 [Antoniadis, Bachas, Fabre,
 Partouche, Taylor]

$\mathcal{A}, \mathcal{M}, \mathcal{K}$
 $\mathcal{N} = 2$

Coefficients determined by string theory

Conjecture for Kähler potential of dilaton and untwisted moduli for T^6/\mathbb{Z}'_6 (up to 1-loop):

$$\begin{aligned}
K = & -\ln(T_0 - \bar{T}_0) - \ln[(T_1 - \bar{T}_1)(T_2 - \bar{T}_2)(T_3 - \bar{T}_3)] - \ln(U_2 - \bar{U}_2) \\
& + c_1 \chi \zeta(3) \sqrt{\frac{(T_1 - \bar{T}_1)(T_2 - \bar{T}_2)(T_3 - \bar{T}_3)}{(T_0 - \bar{T}_0)^3}} \quad \xleftarrow{\textcolor{red}{S^2}} \\
& + c_2 \frac{E_2(U_2)}{(T_2 - \bar{T}_2)(T_0 - \bar{T}_0)} + c_3 \frac{E_2(-1/U_2)}{(T_1 - \bar{T}_1)(T_3 - \bar{T}_3)} + c_4 \frac{E_2(U_3)}{(T_3 - \bar{T}_3)(T_0 - \bar{T}_0)} \\
& + c_5 \frac{1}{\sqrt{(T_1 - \bar{T}_1)(T_2 - \bar{T}_2)(T_3 - \bar{T}_3)(T_0 - \bar{T}_0)}} \quad \xrightarrow{\mathcal{A}, \mathcal{M}, \mathcal{K}} \quad \mathcal{N} = 2
\end{aligned}$$

$\mathcal{T}, \mathcal{A}, \mathcal{M}, \mathcal{K}$
 $\mathcal{N} = 1$

Conjecture for Kähler potential of dilaton and untwisted moduli for T^6/\mathbb{Z}'_6 (up to 1-loop):

$$\begin{aligned}
K = & -\ln(T_0 - \bar{T}_0) - \ln[(T_1 - \bar{T}_1)(T_2 - \bar{T}_2)(T_3 - \bar{T}_3)] - \ln(U_2 - \bar{U}_2) \\
& + c_1 \chi \zeta(3) \sqrt{\frac{(T_1 - \bar{T}_1)(T_2 - \bar{T}_2)(T_3 - \bar{T}_3)}{(T_0 - \bar{T}_0)^3}} \quad \xleftarrow{\textcolor{red}{S^2}} \\
& + c_2 \frac{E_2(U_2)}{(T_2 - \bar{T}_2)(T_0 - \bar{T}_0)} + c_3 \frac{E_2(-1/U_2)}{(T_1 - \bar{T}_1)(T_3 - \bar{T}_3)} + c_4 \frac{E_2(U_3)}{(T_3 - \bar{T}_3)(T_0 - \bar{T}_0)} \\
& + c_5 \frac{1}{\sqrt{(T_1 - \bar{T}_1)(T_2 - \bar{T}_2)(T_3 - \bar{T}_3)(T_0 - \bar{T}_0)}} \quad \xrightarrow{\mathcal{A}, \mathcal{M}, \mathcal{K}} \quad \mathcal{N} = 2
\end{aligned}$$

$\mathcal{T}, \mathcal{A}, \mathcal{M}, \mathcal{K}$
 $\mathcal{N} = 1$

Note:

$$K^{(1)} = 8\mathcal{V}_1^2 G_{\mathcal{V}_1 \mathcal{V}_1}^{(1)}$$

[M.H., Kang]

(follows from relations
between different
metric components)

Outlook

- Work out coefficients in K and field redefinitions
- Additional field redefinitions from α' -corrections?
[Antoniadis, Minasian, Theisen, Vanhove; Grimm, Savelli, Weissenbacher]
- Check correction to EH-term at disk level
[Green, Rudra]
- Applications to string model building?

Хвала!



All the best for the future!